

BONZI

Bonzi is the most reliable plant growth regulator, helping growers maximise production in a consistent and cost-effective way.

Active ingredient

4g/l Paclobutrazol

MAPP 17576

Formulation SC (Suspension Concentrate)

Pack size 1 litre

Efficient


- Fast acting within 3-5 days
- Long lasting minimum 3 weeks
- Improves uniformity of shape & size

Flexible

- Can be applied along the crop cycle
- Can be tank mixed and alternated with other growth regulators
- Can be used in most P&B crops

Reliable

- Works in all circumstances
- Safe to crops
- Delivers robust and compact plants



Bonzi[®]

Ornamental growth regulator

syngenta

Syngenta UK Ltd,
CPC4 Capital Park, Fulbourn, Cambridge, CB21 5XE
01223 883400
www.syngentaornamentals.co.uk

@Syngenta_ornamentals
 Syngenta Ornamentals UK
 Syngenta Ornamentals EAME
 @SyngentaOCUK

Registered in England No. 849037

Mode of action


Bonzi inhibits the production of gibberellins at an early stage of the process, resulting in shorter internodes and more controlled growth. This way Bonzi can manage the growth and shape of a plant which results in a more compact and firm plant that fits with the requirements of the market.



Growth control with Bonzi Plant Growth Regulator Spray Treatment on Calliope Dark Red Geraniums

Uptake and translocation


Bonzi can easily be taken up by the plant through leaves, roots and stems. It is transported upwards to the growing points through the xylem. The distribution is limited within the leaf, therefore an equal coverage of the leaves is critical.



How to use Bonzi in the optimal way

- Avoid tankmixing Bonzi with fungicides or insecticides, as this can influence the effect of Bonzi.
- Apply on a dry crop.
- Ensure that the crop has enough water.
- Apply the appropriate dosage to avoid overdosing (better to spray several times with a lower dose rate than one or two times overdosed).
- After applying Bonzi wait for the crop to dry before overhead watering.
- Ensure the optimum water volume to prevent Bonzi on the ground.
- Do not re-use growing media which are used with treated plants.
- Bonzi has no negative effects on flowers and flower development.
- Earlier flowering after application of Bonzi has been observed in Petunia, Verbena, Fuchsia and Dianthus.
- In Poinsettia Bonzi should be applied before the first week of floral indication to prevent effect on the bract size.

Bonzi in pot & bedding plants



The effect of Bonzi in Poinsettia at different dose rates

syngenta®